

Academy of Sciences of Ukraine
Institute for Condensed Matter Physics

Preprint
ICMP-93-22E

O.Derzhko, A.Moina

Statistical mechanics of one-dimensional $s = \frac{1}{2}$
anisotropic XY model in transverse field with
Dzyaloshinskii-Moriya interaction

L'viv-1994

O.Derzhko, A.Moina

Statistical mechanics of one-dimensional $s = \frac{1}{2}$ anisotropic XY model in transverse field with Dzyaloshinskii-Moriya interaction

For 1D $s = \frac{1}{2}$ anisotropic XY model in transverse field with Dzyaloshinskii-Moriya interaction using Jordan-Wigner transformation the thermodynamical functions, static spin correlation functions, transverse dynamical spin correlation function and connected with it transverse dynamical susceptibility have been obtained. It has been shown that Dzyaloshinskii-Moriya interaction essentially influences the calculated quantities.

©1994 Institute for Condensed Matter Physics

1 Introduction

In 1961 E.Lieb, T.Schultz and D.Mattis in ref. [1] pointed out one type of exactly solvable models of statistical mechanics that is so called 1D $s = \frac{1}{2}$ XY models. Rewriting the Hamiltonian of such chain

$$H = \sum_j \left[(1 + \gamma) s_j^x s_{j+1}^x + (1 - \gamma) s_j^y s_{j+1}^y \right], \quad -1 \leq \gamma \leq 1, \quad (1)$$

$$[s_j^\alpha, s_m^\beta] = i\delta_{jm} s_m^\gamma, \quad \alpha, \beta, \gamma = x, y, z + \text{cyclic permutations} \quad (2)$$

with the help of the raising and lowering operators $s_j^\pm \equiv s_j^x \pm i s_j^y$ in the form

$$H = \frac{1}{2} \sum_j \left[(\gamma s_j^+ s_{j+1}^+ + s_j^+ s_{j+1}^-) + h.c. \right], \quad (3)$$

they noted that the difficulty of diagonalization of the obtained quadratic in operators s^+, s^- form (3) is connected with the commutation rules that these operators satisfy, namely, $[s_j^-, s_m^+] = \delta_{jm} (1 - 2s_m^+ s_m^-)$. Really, they are similar to Fermi-type commutation rules for operators at the same site and to Bose-type commutation rules for operators attached to different sites

$$\{s_j^-, s_j^+\} = 1, \quad (s_j^+)^2 = (s_j^-)^2 = 0;$$

$$[s_j^-, s_m^+] = [s_j^+, s_m^+] = [s_j^-, s_m^-] = 0, \quad j \neq m. \quad (4)$$

That is why one should perform at first Jordan-Wigner transformation (see, besides ref. [1], also refs.[2-4])

$$\begin{aligned} c_1 &= s_1^-, \quad c_j = s_j^- P_{j-1} = P_{j-1} s_j^-, \quad j = 2, \dots, N, \\ c_1^+ &= s_1^+, \quad c_j^+ = s_j^+ P_{j-1} = P_{j-1} s_j^+, \quad j = 2, \dots, N, \end{aligned} \quad (5)$$

where Jordan-Wigner factor is denoted by $P_j \equiv \prod_{n=1}^j (-2s_n^z)$. The introduced operators really obey Fermi commutation rules. From (5) it follows that

$$c_j^+ c_j = s_j^+ P_{j-1}^2 s_j^- = s_j^+ s_j^-, \quad c_j c_j^+ = s_j^- s_j^+, \quad c_j^+ c_j^+ = s_j^+ s_j^+, \quad c_j c_j = s_j^- s_j^-, \quad (6)$$

since $P_j^2 = \prod_{n=1}^j (-2s_n^z)^2 = \prod_{n=1}^j 4(s_n^z)^2 = 1$, and the commutation rules at the same site remain of Fermi-type. Consider then a product of c -operators at different sites

$$c_n^+ c_m = s_n^+ \prod_{p=1}^{n-1} (-2s_p^z) \prod_{j=1}^{m-1} (-2s_j^z) s_m^- = s_n^+ \prod_{j=n}^{m-1} (-2s_j^z) s_m^-, \quad (7)$$

putting here for definiteness $n < m$. Since $s_j^\pm(-2s_j^z) = \pm s_j^\pm$ and $(-2s_j^z)s_j^\pm = \mp s_j^\pm$, and consequently

$$c_m c_n^+ = s_m^- \prod_{j=n}^{m-1} (-2s_j^z) s_n^+ = -s_n^+ \prod_{j=n}^{m-1} (-2s_j^z) s_m^-, \quad (8)$$

one gets $c_n^+ c_m = -c_m c_n^+$. Similarly one finds that $c_n^+ c_m^+ = -c_m^+ c_n^+$, $c_n c_m = -c_m c_n$. Thus the introduced in (5) operators are Fermi-type operators

$$\{c_j, c_l^+\} = \delta_{jl}, \quad \{c_j^+, c_l^+\} = \{c_j, c_l\} = 0. \quad (9)$$

Since $P_j^2 = 1$, $P_j = \exp(\pm i\pi \sum_{n=1}^j s_n^+ s_n^-)$ (because $\exp[\pm i\pi \sum_{n=1}^j (\frac{1}{2} + s_n^z)] = \prod_{n=1}^j (-2s_n^z)$), $s_j^+ s_j^- = c_j^+ c_j$, it is easy to write the inverse to (5) transformation

$$\begin{aligned} s_1^- &= c_1, \quad s_j^- = c_j \exp(\pm i\pi \sum_{n=1}^{j-1} c_n^+ c_n) = \exp(\pm i\pi \sum_{n=1}^{j-1} c_n^+ c_n) c_j, \quad j = 2, \dots, N, \\ s_1^+ &= c_1^+, \quad s_j^+ = c_j^+ \exp(\pm i\pi \sum_{n=1}^{j-1} c_n^+ c_n) = \exp(\pm i\pi \sum_{n=1}^{j-1} c_n^+ c_n) c_j^+, \quad j = 2, \dots, N. \end{aligned} \quad (10)$$

Returning to the Hamiltonian (3) one notes that the products of two Pauli operators at neighbouring sites transform into such products of Fermi operators

$$\begin{aligned} c_j^+ c_{j+1}^+ &= s_j^+ (-2s_j^z) s_{j+1}^+ = s_j^+ s_{j+1}^+, \\ c_j^+ c_{j+1} &= s_j^+ (-2s_j^z) s_{j+1}^- = s_j^+ s_{j+1}^-, \\ c_j c_{j+1}^+ &= s_j^- (-2s_j^z) s_{j+1}^+ = -s_j^- s_{j+1}^+, \\ c_j c_{j+1} &= s_j^- (-2s_j^z) s_{j+1}^- = -s_j^- s_{j+1}^-. \end{aligned} \quad (11)$$

Usually bearing in mind the study of thermodynamical properties of the system that requires the performance of thermodynamical limit $N \rightarrow \infty$, the periodic boundary conditions are implied

$$s_{N+1}^\alpha \equiv s_1^\alpha, \quad \alpha = x, y, z. \quad (12)$$

In connection with this in (3) there are products of the form

$$\begin{aligned} s_N^+ s_{N+1}^+ &= s_N^+ s_1^+ = s_1^+ s_N^+ = c_1^+ c_N^+ \prod_{p=1}^{N-1} (-2s_p^z) = c_1^+ c_N^+ P, \quad P \equiv P_N, \\ s_N^+ s_{N+1}^- &= c_1 c_N^+ P, \quad s_N^- s_{N+1}^+ = -c_1^+ c_N P, \quad s_N^- s_{N+1}^- = -c_1 c_N P. \end{aligned} \quad (13)$$

Gathering the obtained terms one finds that for the ring

$$H = H^- + BP^+ = H^+P^+ + H^-P^-. \quad (14)$$

Here

$$H^\pm \equiv \frac{1}{2} \sum_{j=1}^N \left[(\gamma c_j^\pm c_{j+1}^\pm + c_j^\pm c_{j+1}) + h.c. \right], \quad (15)$$

the difference between H^+ and H^- is only in the implied boundary conditions: for H^+ they are antiperiodic

$$c_j^+ = -c_{j+N}^+, \quad c_j = -c_{j+N}, \quad (16)$$

and for H^- they are periodic

$$c_j^+ = c_{j+N}^+, \quad c_j = c_{j+N}; \quad (17)$$

$B \equiv H^+ - H^- = - \left[(\gamma c_N^+ c_1^+ + c_N^+ c_1) + h.c. \right]$ is the boundary term; $P^\pm \equiv (1 \pm P)/2$ are orthogonal projectors ($P^+ + P^- = 1$, $(P^\pm)^2 = P^\pm$, $P^\pm P^\mp = 0$), besides this $[H^\pm, P] = [H^\pm, P^\pm] = 0$. For the open chain with free ends (then in the sum in (1) the range of the summation index is $j = 1, \dots, N-1$) the Hamiltonian after fermionization has the form

$$H = \frac{1}{2} \sum_{j=1}^{N-1} \left[(\gamma c_j^+ c_{j+1}^+ + c_j^+ c_{j+1}) + h.c. \right]. \quad (18)$$

Formulae (14), (15) or (18) realize the reformulation of the initial Hamiltonian (1) in terms of fermions. They are the starting point for further study of statistical mechanics of models like (1). Besides it appears [5,6] that for calculation of free energy

$$f \equiv -\frac{1}{\beta} \lim_{N \rightarrow \infty} \left[\frac{1}{N} \text{Sp} \exp(-\beta H) \right] \quad (19)$$

or static spin correlation functions

$$\langle s_{j_1}^{\alpha_1} \dots s_{j_n}^{\alpha_n} \rangle \equiv \lim_{N \rightarrow \infty} \left\{ \text{Sp} \left[\exp(-\beta H) s_{j_1}^{\alpha_1} \dots s_{j_n}^{\alpha_n} \right] / \text{Sp} \exp(-\beta H) \right\} \quad (20)$$

the boundary term may be omitted and hence one has to consider a system of free fermions. It is more difficult to calculate the dynamical correlation functions. Really,

$$\begin{aligned} s_j^z(t) &\equiv \exp(\imath H t) s_j^z \exp(-\imath H t) = \\ &= \exp(\imath H^+ t) s_j^z \exp(-\imath H^+ t) P^+ + \exp(\imath H^- t) s_j^z \exp(-\imath H^- t) P^- = \\ &= P^+ \exp(\imath H^+ t) s_j^z \exp(-\imath H^+ t) + P^- \exp(\imath H^- t) s_j^z \exp(-\imath H^- t) \end{aligned} \quad (21)$$

(owing to the following relation that is valid for arbitrary function of $H = H^+P^+ + H^-P^-$: $f(H) \equiv \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} (H^+P^+ + H^-P^-)^n = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} \times \times [(H^+)^n P^+ + (H^-)^n P^-] = f(H^+)P^+ + f(H^-)P^- = P^+f(H^+) + P^-f(H^-)$) in contrast to

$$\begin{aligned} s_j^{x,y}(t) &= \\ &= \exp(\imath H^+ t) s_j^{x,y} \exp(-\imath H^- t) P^- + \exp(\imath H^- t) s_j^{x,y} \exp(-\imath H^+ t) P^+ = \\ &= P^+ \exp(\imath H^+ t) s_j^{x,y} \exp(-\imath H^- t) + P^- \exp(\imath H^- t) s_j^{x,y} \exp(-\imath H^+ t). \end{aligned} \quad (22)$$

In accordance with (21) the pair transverse correlation function in the thermodynamical limit can be written as

$$\langle s_j^z(t) s_{j+n}^z \rangle = \frac{Sp \left[\exp(-\beta H^-) \exp(\imath H^- t) s_j^z \exp(-\imath H^- t) s_{j+n}^z \right]}{Sp \exp(-\beta H^-)} \quad (23)$$

and hence may be calculated with c -cyclic Hamiltonian. Whereas the pair longitudinal correlation function in accordance with (22) in the thermodynamical limit can be written as

$$\langle s_j^x(t) s_{j+n}^x \rangle = \frac{Sp \left[\exp(-\beta H^-) \exp(\imath H^- t) s_j^x \exp(-\imath H^- t) O^-(t) s_{j+n}^x \right]}{Sp \exp(-\beta H^-)} \quad (24)$$

where $O^-(t) \equiv \exp(\imath H^- t) \exp(-\imath(H^- + B)t)$. The calculation with c -cyclic Hamiltonian that neglects the boundary term B yields the approximate result that, in particular, is incorrect in the limit of Ising model ($\gamma = 1$) (see, for instance, [7]). It is interesting to note that the calculation of the four-spin correlation function in the thermodynamical limit involves only c -cyclic Hamiltonian

$$\begin{aligned} \langle s_{j_1}^x(t) s_{j_2}^x(t) s_{j_3}^x s_{j_4}^x \rangle &= \\ &= \frac{Sp \left[\exp(-\beta H^-) \exp(\imath H^- t) s_{j_1}^x s_{j_2}^x \exp(-\imath H^- t) s_{j_3}^x s_{j_4}^x \right]}{Sp \exp(-\beta H^-)}. \end{aligned} \quad (25)$$

Thus here as in the case (23) one comes to calculation of the dynamical correlation functions of the system of non-interacting fermions (see [8]). The calculation of the pair longitudinal correlation function, in spite of a great number of papers dealing with this problem, remains an open point of statistical mechanics of 1D $s = \frac{1}{2}$ XY models. Among other interesting and principal questions of the theory of 1D $s = \frac{1}{2}$ XY models one may mention

the investigation of nonequilibrium properties of such models (see, for example, [9]) and the examination of the properties of disordered versions of such models (see, for example, [10]).

It is necessary to stress the essential features of the present consideration:

- the dimension of space $D=1$;
- the value of spin $s = \frac{1}{2}$;
- interactions occur only between neighbouring spins (otherwise the Hamiltonian will contain the terms that are the products of more than two Fermi operators);
- only x and y components of spins interact and the field that may be included should be transverse (the interaction of z components, for instance, leads to the appearance in the Hamiltonian of the terms that are the products of four Fermi operators).

In connection with this it is easy to point out the model that has more general than in (1) form of interspin interaction, and that still allows the described consideration. Really, considering the additional terms in the Hamiltonian that have form $\sum_j (J^{xy} s_j^x s_{j+1}^y + J^{yx} s_j^y s_{j+1}^x)$ one notes that after fermionization they do not change the form of the Hamiltonian (14), (15) or (18), and lead only to changes in the values of constants. The Hamiltonian of the generalized 1D $s = \frac{1}{2}$ anisotropic XY model in transverse field that as a matter of fact will be studied in the present paper is given by

$$H = \Omega \sum_j s_j^z + \sum_j (J^{xx} s_j^x s_{j+1}^x + J^{xy} s_j^x s_{j+1}^y + J^{yx} s_j^y s_{j+1}^x + J^{yy} s_j^y s_{j+1}^y). \quad (26)$$

Before starting the examination of this model it is worthwhile to mention its possible physical application [11]. For this purpose let's perform the transformation of rotation around axis z over an angle α

$$\begin{aligned} \tilde{s}_j^x &= s_j^x \cos \alpha + s_j^y \sin \alpha, & \tilde{s}_j^y &= -s_j^x \sin \alpha + s_j^y \cos \alpha, & \tilde{s}_j^z &= s_j^z; \\ s_j^x &= \tilde{s}_j^x \cos \alpha - \tilde{s}_j^y \sin \alpha, & s_j^y &= \tilde{s}_j^x \sin \alpha + \tilde{s}_j^y \cos \alpha, & s_j^z &= \tilde{s}_j^z. \end{aligned} \quad (27)$$

Then rewriting at first new terms in sum in the Hamiltonian (26) in the form

$$\frac{J^{xy} + J^{yx}}{2} (s_j^x s_{j+1}^y + s_j^y s_{j+1}^x) + \frac{J^{xy} - J^{yx}}{2} (s_j^x s_{j+1}^y - s_j^y s_{j+1}^x), \quad (28)$$

taking into account that the terms $(s_j^x s_{j+1}^y - s_j^y s_{j+1}^x)$ are invariant under transformation (27) and

$$\begin{aligned}
& J^{xx} s_j^x s_{j+1}^x + \frac{J^{xy} + J^{yx}}{2} (s_j^x s_{j+1}^y + s_j^y s_{j+1}^x) + J^{yy} s_j^y s_{j+1}^y = \\
& = \left(J^{xx} \cos^2 \alpha + \frac{J^{xy} + J^{yx}}{2} \sin 2\alpha + J^{yy} \sin^2 \alpha \right) \tilde{s}_j^x \tilde{s}_{j+1}^x + \\
& + \left(\frac{J^{yy} - J^{xx}}{2} \sin 2\alpha + \frac{J^{xy} + J^{yx}}{2} \cos 2\alpha \right) (\tilde{s}_j^x \tilde{s}_{j+1}^y + \tilde{s}_j^y \tilde{s}_{j+1}^x) + \\
& + \left(J^{xx} \sin^2 \alpha - \frac{J^{xy} + J^{yx}}{2} \sin 2\alpha + J^{yy} \cos^2 \alpha \right) \tilde{s}_j^y \tilde{s}_{j+1}^y, \quad (29)
\end{aligned}$$

and choosing the parameter of transformation α from the condition $(J^{xy} + J^{yx}) \cos 2\alpha - (J^{xx} - J^{yy}) \sin 2\alpha = 0$, one will have

$$H = \Omega \sum_j \tilde{s}_j^z + \sum_j \left[J^x \tilde{s}_j^x \tilde{s}_{j+1}^x + J^y \tilde{s}_j^y \tilde{s}_{j+1}^y + D(\tilde{s}_j^x \tilde{s}_{j+1}^y - \tilde{s}_j^y \tilde{s}_{j+1}^x) \right], \quad (30)$$

where

$$\begin{aligned}
J^x & \equiv J^{xx} \cos^2 \alpha + \frac{J^{xy} + J^{yx}}{2} \sin 2\alpha + J^{yy} \sin^2 \alpha, \\
J^y & \equiv J^{xx} \sin^2 \alpha - \frac{J^{xy} + J^{yx}}{2} \sin 2\alpha + J^{yy} \cos^2 \alpha, \\
D & \equiv \frac{J^{xy} - J^{yx}}{2}, \quad \tan 2\alpha = \frac{J^{xy} + J^{yx}}{J^{xx} - J^{yy}}. \quad (31)
\end{aligned}$$

In the term that is proportional to D one easily recognizes z component of the vector $[\vec{s}_j \times \vec{s}_{j+1}]$ that is the so called Dzyaloshinskii-Moriya interaction. It was at first introduced phenomenologically by I.E. Dzyaloshinskii [12] and then derived by T.Moriya [13] by extending Anderson's theory of superexchange interactions [14] to include spin-orbital coupling (see, for example, ref.[15]). The model with relativistic Dzyaloshinskii-Moriya interaction together with ANNNI model are widely used in microscopic theory of crystals with incommensurate phase [16,17]. In the classical case Dzyaloshinskii-Moriya interaction may lead to the appearance of the spiral spin structure. The possibility of the appearance of spiral structure in quantum case has been studied in ref.[11] where for this purpose pair static spin correlation functions have been estimated.

Except the mentioned paper [11] the problem of statistical mechanics of 1D $s = \frac{1}{2}$ XY type model with the Hamiltonian (26) or (30) as far as the

authors know was not considered yet ¹. At the present paper an attempt to fill up this gap by the generalization for this case of the well-known scheme of consideration of 1D $s = \frac{1}{2}$ XY model has been made. In section 2 the transformation of the Hamiltonian to the initial form for further examination of statistical properties is presented. In section 3 the thermodynamical properties of the model are considered, and in section 4 it is shown how to calculate the static spin correlation functions in this model. The dynamics of transverse spin correlations and the transverse dynamical susceptibility are studied in section 5. The conclusions form section 6.

2 Transformation of the Hamiltonian

In the spirit of above described approach the Hamiltonian of the model (26) at first should be rewritten with the help of the raising and lowering operators in the form that is similar to (3)

$$\begin{aligned}
H = & \Omega \sum_{j=1}^N \left(s_j^+ s_j^- - \frac{1}{2} \right) + \\
& + \sum_{j=1}^N \left(J^{++} s_j^+ s_{j+1}^+ + J^{+-} s_j^+ s_{j+1}^- + J^{-+} s_j^- s_{j+1}^+ + J^{--} s_j^- s_{j+1}^- \right), \\
J^{++} \equiv & [J^{xx} - J^{yy} - \imath(J^{xy} + J^{yx})] / 4 = (J^{--})^*, \\
J^{+-} \equiv & [J^{xx} + J^{yy} + \imath(J^{xy} - J^{yx})] / 4 = (J^{-+})^*; \tag{32}
\end{aligned}$$

here the periodic boundary conditions (12) are imposed. The Hamiltonian of the model (32) after Jordan-Wigner transformation (5), (10) will have the form that is similar to (14), (15)

$$\begin{aligned}
H = & H^+ P^+ + H^- P^-, \\
H^\pm \equiv & \Omega \sum_{j=1}^N \left(c_j^+ c_j^- - \frac{1}{2} \right) + \sum_{j=1}^N \left(J^{++} c_j^+ c_{j+1}^+ + J^{+-} c_j^+ c_{j+1}^- - \right. \\
& \left. - J^{-+} c_j^- c_{j+1}^+ - J^{--} c_j^- c_{j+1}^- \right); \tag{33}
\end{aligned}$$

besides H^- is c -cyclic and H^+ is c -anticyclic quadratic forms in Fermi operators. After Fourier transformation

$$c_\kappa^+ = \frac{1}{\sqrt{N}} \sum_{j=1}^N e^{-\imath \kappa j} c_j^+, \quad c_\kappa = \frac{1}{\sqrt{N}} \sum_{j=1}^N e^{\imath \kappa j} c_j,$$

¹In ref.[18] on the base of the model with Hamiltonian (26) the problem about the validity of the Bose commutation rules approximation for spin operators has been examined.

$$c_j^+ = \frac{1}{\sqrt{N}} \sum_{\kappa} e^{i\kappa j} c_{\kappa}^+, \quad c_j = \frac{1}{\sqrt{N}} \sum_{\kappa} e^{-i\kappa j} c_{\kappa}, \quad (34)$$

with $\kappa = \kappa^- \equiv 2\pi n/N$ for H^- and $\kappa = \kappa^+ \equiv 2\pi(n + 1/2)/N$ for H^+ , $n = -N/2, -N/2 + 1, \dots, N/2 - 1$ (for even N), $n = -(N - 1)/2, -(N - 1)/2 + 1, \dots, (N - 1)/2$ (for odd N) $H^{\pm}$ can be rewritten in the form

$$H^{\pm} = \sum_{\kappa} \left[-\frac{\Omega}{2} + \epsilon_{\kappa} c_{\kappa}^+ c_{\kappa} - i \sin \kappa \left(J^{++} c_{\kappa}^+ c_{-\kappa}^+ + J^{--} c_{\kappa} c_{-\kappa} \right) \right],$$

$$\epsilon_{\kappa} \equiv \epsilon_{\kappa}^{(+)} + \epsilon_{\kappa}^{(-)}, \quad \epsilon_{\kappa}^{(+)} \equiv \Omega + \frac{J^{xx} + J^{yy}}{2} \cos \kappa, \quad \epsilon_{\kappa}^{(-)} \equiv \frac{J^{xy} - J^{yx}}{2} \sin \kappa \quad (35)$$

(here the following relations

$$\sum_{\kappa} e^{-i\kappa} c_{\kappa}^+ c_{-\kappa}^+ = -i \sum_{\kappa} \sin \kappa c_{\kappa}^+ c_{-\kappa}^+, \quad \sum_{\kappa} e^{i\kappa} c_{\kappa} c_{-\kappa} = i \sum_{\kappa} \sin \kappa c_{\kappa} c_{-\kappa} \quad (36)$$

were used). The diagonalization of the quadratic forms is finished up by the Bogolyubov transformation

$$\begin{aligned} \beta_{\kappa} &= x_{\kappa} c_{\kappa} + y_{\kappa} c_{-\kappa}^+, & \beta_{-\kappa}^+ &= y_{-\kappa}^* c_{\kappa} + x_{-\kappa}^* c_{-\kappa}^+, \\ c_{\kappa} &= \left(-x_{-\kappa}^* \beta_{\kappa} + y_{\kappa} \beta_{-\kappa}^+ \right) / \Delta_{\kappa}, & c_{-\kappa}^+ &= \left(y_{-\kappa}^* \beta_{\kappa} - x_{\kappa} \beta_{-\kappa}^+ \right) / \Delta_{\kappa}, \\ \Delta_{\kappa} &\equiv y_{\kappa} y_{-\kappa}^* - x_{\kappa} x_{-\kappa}^* \neq 0. \end{aligned} \quad (37)$$

β -operators remain of Fermi type if

$$|x_{\kappa}|^2 + |y_{\kappa}|^2 = 1, \quad \frac{x_{\kappa}}{y_{\kappa}} + \frac{x_{-\kappa}}{y_{-\kappa}} = 0. \quad (38)$$

The transformed Hamiltonian contains the operator terms proportional only to $\beta_{\kappa}^+ \beta_{\kappa}$ if

$$\epsilon_{\kappa}^{(+)} + i \sin \kappa \left(J^{++} \frac{x_{\kappa}}{y_{\kappa}} - J^{--} \frac{y_{-\kappa}}{x_{-\kappa}} \right) = 0. \quad (39)$$

The condition (39) and the second condition in (38) yield

$$\begin{aligned} \frac{x_{\kappa}}{y_{\kappa}} &= i \frac{\epsilon_{\kappa}^{(+)} \mp \mathcal{E}_{\kappa}}{2|J^{++}| \sin \kappa} \exp(-i \arg J^{++}), \\ \mathcal{E}_{\kappa} &\equiv \sqrt{(\epsilon_{\kappa}^{(+)})^2 + 4|J^{++}|^2 \sin^2 \kappa}. \end{aligned} \quad (40)$$

Taking into account the first condition in (38) one finds that for lower sign in (40)

$$\begin{aligned} x_{\kappa} &= 2i|J^{++}| \sin \kappa \exp(-i \arg J^{++}) / \sqrt{2\mathcal{E}_{\kappa}(\mathcal{E}_{\kappa} - \epsilon_{\kappa}^{(+)})}, \\ y_{\kappa} &= \sqrt{(\mathcal{E}_{\kappa} - \epsilon_{\kappa}^{(+)}) / 2\mathcal{E}_{\kappa}}, \end{aligned} \quad (41)$$

besides $\Delta_\kappa = 1$, $E_\kappa = \epsilon_\kappa^{(-)} + \mathcal{E}_\kappa$. For upper sign in (40)

$$\begin{aligned} x_\kappa &= -2l|J^{++}| \sin \kappa \exp(-i \arg J^{++}) / \sqrt{2\mathcal{E}_\kappa(\mathcal{E}_\kappa + \epsilon_\kappa^{(+)})}, \\ y_\kappa &= \sqrt{(\mathcal{E}_\kappa + \epsilon_\kappa^{(+)})/2\mathcal{E}_\kappa}; \end{aligned} \quad (42)$$

besides $\Delta_\kappa = 1$, $E_\kappa = \epsilon_\kappa^{(-)} - \mathcal{E}_\kappa$. Thus in a result of Bogolyubov transformation (37) one gets

$$H^\pm = \sum_\kappa E_\kappa \left(\beta_\kappa^+ \beta_\kappa - 1/2 \right), \quad \{\beta_\kappa, \beta_\pi^+\} = \delta_{\kappa\pi}, \quad \{\beta_\kappa, \beta_\pi\} = \{\beta_\kappa^+, \beta_\pi^+\} = 0. \quad (43)$$

It is important to note that in contrast to anisotropic XY model because of inequality $J^{xy} \neq J^{yx}$ one has $E_\kappa \neq E_{-\kappa}$. This is connected with the absence of symmetry with respect to spatial inversion. Really, the Hamiltonian of the model (26) $H(\Omega, J^{xx}, J^{xy}, J^{yx}, J^{yy})$ under the action of spatial inversion, that leads to change of indexes j to $-j$ or $N-j$, $j+1$ to $N-j-1$, transforms into $H(\Omega, J^{xx}, J^{yx}, J^{xy}, J^{yy})$.

In fig. 1 it is shown how the presence of Dzyaloshinskii-Moriya interaction influences the dependence of $E_\kappa = \epsilon_\kappa^{(-)} + \mathcal{E}_\kappa$ on κ in de Gennes model (1 : $D = 0, \quad \Omega = 0, \quad 1' : D = J^{xx}, \quad \Omega = 0; \quad 2 : D = 0, \quad \Omega = J^{xx}, \quad 2' : D = J^{xx}, \quad \Omega = J^{xx}$). In fig. 2 the same is depicted for the case of isotropic XY model.

It is worthwhile to note that the spectrum of elementary excitations in the model under consideration as it follows from the expression for ground state energy (46) is given by $|E_\kappa|$.

3 Thermodynamics

For investigation of thermodynamical properties of the model in question let's calculate the free energy per site in the limit $N \rightarrow \infty$. In accordance with refs.[5,6] for such calculation one can use c -cyclic Hamiltonian and thus

$$f = -\frac{1}{\beta} \lim_{N \rightarrow \infty} \left[\frac{1}{N} \ln Sp \exp(-\beta H^-) \right]. \quad (44)$$

The diagonalized quadratic in Fermi operators form H^- involved in (44) has the form (43), and owing to this one easily obtains the desired result

$$f = -\frac{1}{\beta} \lim_{N \rightarrow \infty} \left\{ \frac{1}{N} \ln Sp \exp \left[-\beta \sum_\kappa E_\kappa \left(\beta_\kappa^+ \beta_\kappa - \frac{1}{2} \right) \right] \right\} =$$

$$\begin{aligned}
&= -\frac{1}{\beta} \lim_{N \rightarrow \infty} \left\{ \frac{1}{N} \ln Sp \prod_{\kappa} \exp \left[-\beta E_{\kappa} \left(\beta_{\kappa}^+ \beta_{\kappa} - \frac{1}{2} \right) \right] \right\} = \\
&= -\frac{1}{\beta} \lim_{N \rightarrow \infty} \left[\frac{1}{N} \ln \left(\prod_{\kappa} 2 \cosh \frac{\beta E_{\kappa}}{2} \right) \right] = -\frac{1}{\beta} \lim_{N \rightarrow \infty} \left[\frac{1}{N} \sum_{\kappa} \ln \left(2 \cosh \frac{\beta E_{\kappa}}{2} \right) \right] = \\
&= -\frac{1}{\beta} \frac{1}{2\pi} \int_{-\pi}^{\pi} d\kappa \ln \left(2 \cosh \frac{\beta E_{\kappa}}{2} \right). \tag{45}
\end{aligned}$$

Knowing the free energy (45) one finds the energy of the ground state

$$e = \lim_{\beta \rightarrow \infty} f = -\frac{1}{4\pi} \int_{-\pi}^{\pi} d\kappa |E_{\kappa}|, \tag{46}$$

the entropy

$$s = \beta^2 \frac{\partial f}{\partial \beta} = \frac{1}{2\pi} \int_{-\pi}^{\pi} d\kappa \ln \left(2 \cosh \frac{\beta E_{\kappa}}{2} \right) - \frac{1}{2\pi} \int_{-\pi}^{\pi} d\kappa \frac{\beta E_{\kappa}}{2} \tanh \frac{\beta E_{\kappa}}{2}, \tag{47}$$

the specific heat

$$c = -\beta \frac{\partial s}{\partial \beta} = \frac{1}{2\pi} \int_{-\pi}^{\pi} d\kappa \left(\frac{\beta E_{\kappa}}{2} \right)^2 \left(\cosh \frac{\beta E_{\kappa}}{2} \right)^{-2}, \tag{48}$$

the transverse magnetization

$$\langle \frac{1}{N} \sum_{j=1}^N s_j^z \rangle = \frac{\partial f}{\partial \Omega} = -\frac{1}{4\pi} \int_{-\pi}^{\pi} d\kappa \frac{\partial E_{\kappa}}{\partial \Omega} \tanh \frac{\beta E_{\kappa}}{2}, \tag{49}$$

the static transverse susceptibility

$$\begin{aligned}
\chi_{zz} &= \frac{\partial \langle \frac{1}{N} \sum_{j=1}^N s_j^z \rangle}{\partial \Omega} = \\
&= -\frac{1}{4\pi} \int_{-\pi}^{\pi} d\kappa \frac{\partial^2 E_{\kappa}}{\partial \Omega^2} \tanh \frac{\beta E_{\kappa}}{2} - \frac{\beta}{8\pi} \int_{-\pi}^{\pi} d\kappa \left(\frac{\partial E_{\kappa}}{\partial \Omega} \right)^2 \left(\cosh \frac{\beta E_{\kappa}}{2} \right)^{-2}. \tag{50}
\end{aligned}$$

In order to illustrate the influence of the additional interactions on thermodynamical properties let's present the results of numerical calculations of the specific heat (48) as a function of temperature (figs. 3,4 (1: $D = 0$, $\Omega = 0$, $1' : D = J^{xx}$, $\Omega = 0$; 2: $D = 0$, $\Omega = J^{xx}$, $2' : D = J^{xx}$, $\Omega = J^{xx}$)) and of transverse magnetization (49) as a function of transverse field (figs. 5,6 (1: $D = 0$, $\beta = 1/J^{xx}$, $1' : D = J^{xx}$, $\beta = 1/J^{xx}$; 2: $D = 0$, $\beta = 1000/J^{xx}$, $2' : D = J^{xx}$, $\beta = 1000/J^{xx}$)) and of temperature (figs. 7,8 (1: $D = 0$, $\Omega = 0$, $1' : D = J^{xx}$, $\Omega = 0$; 2: $D = 0$, $\Omega = J^{xx}$, $2' : D = J^{xx}$, $\Omega = J^{xx}$)) for de Gennes model and for isotropic XY model in the presence of Dzyaloshinskii-Moriya interaction.

4 Static spin correlation functions

For the investigation of the spin structure in the model under consideration let's introduce the static spin correlation functions. Due to the possibility of their calculation with the help of c -cyclic Hamiltonian the initial formula for their evaluation can be rewritten in the form

$$\langle s_{j_1}^{\alpha_1} \dots s_{j_n}^{\alpha_n} \rangle = \lim_{N \rightarrow \infty} \left\{ Sp \left[\exp(-\beta H^-) s_{j_1}^{\alpha_1} \dots s_{j_n}^{\alpha_n} \right] / Sp \exp(-\beta H^-) \right\}. \quad (51)$$

Let's introduce then φ -operators that owing to (37), (41) are the following linear combinations of β -operators

$$\begin{aligned} \varphi_j^\pm &\equiv c_j^+ \pm c_j = \sum_{\kappa} \left(\lambda_{j\kappa}^\pm \beta_\kappa^+ \pm \mu_{j\kappa}^\pm \beta_{-\kappa} \right), \\ \lambda_{j\kappa}^\pm &\equiv \frac{1}{\sqrt{N}} e^{i\kappa j} (x_\kappa \pm y_\kappa), \quad \mu_{j\kappa}^\pm \equiv \frac{1}{\sqrt{N}} e^{i\kappa j} [x_\kappa \exp(2i \arg J^{++}) \pm y_\kappa], \end{aligned} \quad (52)$$

and in terms of which the spin operators can be presented as

$$s_n^x = \frac{1}{2} \prod_{j=1}^{n-1} (\varphi_j^+ \varphi_j^-) \varphi_n^+, \quad s_n^y = \frac{1}{2i} \prod_{j=1}^{n-1} (\varphi_j^+ \varphi_j^-) \varphi_n^-, \quad s_n^z = \frac{1}{2} \varphi_n^- \varphi_n^+. \quad (53)$$

φ -operators obey the following commutation relations

$$\{\varphi_i^+, \varphi_j^-\} = 0, \quad \{\varphi_i^+, \varphi_j^+\} = 2\delta_{ij}, \quad \{\varphi_i^-, \varphi_j^-\} = -2\delta_{ij}, \quad (54)$$

besides $(\varphi_j^+ \varphi_j^-)^2 = 1$ and

$$[\varphi_i^\pm, \varphi_j^\pm \varphi_j^\mp] = 2\delta_{ij} \varphi_i^\mp, \quad [\varphi_i^+ \varphi_i^-, \varphi_j^\pm \varphi_j^\mp] = 0. \quad (55)$$

That is why the calculation of static spin correlation functions after substitution of (53) into (51) and exploiting of (54), (55) reduces to application of Wick-Bloch-de Dominicis theorem. The theorem states that the mean value of the product of even number of φ operators with the Hamiltonian H^- (43) is equal to the sum of all possible full systems of contractions of this product; if the number of φ operators in the product is odd the mean value of the product is equal to zero. The full system of contractions of the product of even number of Fermi-type operators forms so called Pfaffian the square of which is equal to the determinant of antisymmetric matrix constructed in a certain way from elementary contractions [19,20].

Thus let's consider the calculation of elementary contractions. One has

$$\langle \varphi_j^+ \varphi_{j+n}^+ \rangle = \sum_{\kappa_1, \kappa_2} \left(\lambda_{j\kappa_1}^+ \mu_{j+n, \kappa_2}^+ \langle \beta_{\kappa_1}^+ \beta_{-\kappa_2} \rangle + \mu_{j\kappa_1}^+ \lambda_{j+n, \kappa_2}^+ \langle \beta_{-\kappa_1} \beta_{\kappa_2}^+ \rangle \right) \quad (56)$$

(here evident relations $\langle \beta_{\kappa_1}^+ \beta_{\kappa_2}^+ \rangle = \langle \beta_{-\kappa_1} \beta_{-\kappa_2} \rangle = 0$ were used). Since

$$\begin{aligned} \langle \beta_{\kappa_1}^+ \beta_{-\kappa_2} \rangle &= \delta_{\kappa_1, -\kappa_2} / (1 + e^{\beta E_{\kappa_1}}) = \delta_{\kappa_1, -\kappa_2} f_{\kappa_1}, \\ \langle \beta_{-\kappa_1} \beta_{\kappa_2}^+ \rangle &= \delta_{-\kappa_1, \kappa_2} e^{\beta E_{\kappa_2}} / (1 + e^{\beta E_{\kappa_2}}) = \delta_{-\kappa_1, \kappa_2} e^{\beta E_{\kappa_2}} f_{\kappa_2}, \end{aligned} \quad (57)$$

where $f_{\kappa} \equiv 1/(1 + e^{\beta E_{\kappa}})$ and in accordance with (52)

$$\begin{aligned} \lambda_{j\kappa}^+ \mu_{j+n, -\kappa}^+ &= \frac{1}{N} e^{-i\kappa n} (1 + S_{\kappa}), \quad \mu_{j, -\kappa}^+ \lambda_{j+n, \kappa}^+ = \frac{1}{N} e^{i\kappa n} (1 + S_{\kappa}), \\ S_{\kappa} &\equiv \frac{2|J^{++}| \sin(\arg J^{++}) \sin \kappa}{\mathcal{E}_{\kappa}} \end{aligned} \quad (58)$$

one has

$$\langle \varphi_j^+ \varphi_{j+n} \rangle = \frac{1}{N} \sum_{\kappa} e^{i\kappa n} (1 + S_{\kappa}) - \frac{2i}{N} \sum_{\kappa} \sin(\kappa n) (1 + S_{\kappa}) f_{\kappa}. \quad (59)$$

Similarly one finds that

$$\langle \varphi_j^- \varphi_{j+n}^- \rangle = -\frac{1}{N} \sum_{\kappa} e^{i\kappa n} (1 - S_{\kappa}) + \frac{2i}{N} \sum_{\kappa} \sin(\kappa n) (1 - S_{\kappa}) f_{\kappa} \quad (60)$$

and

$$\begin{aligned} \langle \varphi_j^+ \varphi_{j+n}^- \rangle &= \\ &= \frac{1}{N} \sum_{\kappa} e^{i\kappa n} \left(\frac{\epsilon_{\kappa}^{(+)}}{\mathcal{E}_{\kappa}} + iC_{\kappa} \right) - \frac{2}{N} \sum_{\kappa} \left[\cos(\kappa n) \frac{\epsilon_{\kappa}^{(+)}}{\mathcal{E}_{\kappa}} - \sin(\kappa n) C_{\kappa} \right] f_{\kappa}, \end{aligned} \quad (61)$$

$$\begin{aligned} \langle \varphi_j^- \varphi_{j+n}^+ \rangle &= \\ &= \frac{1}{N} \sum_{\kappa} e^{i\kappa n} \left(\frac{-\epsilon_{\kappa}^{(+)}}{\mathcal{E}_{\kappa}} + iC_{\kappa} \right) + \frac{2}{N} \sum_{\kappa} \left[\cos(\kappa n) \frac{\epsilon_{\kappa}^{(+)}}{\mathcal{E}_{\kappa}} + \sin(\kappa n) C_{\kappa} \right] f_{\kappa}, \\ C_{\kappa} &\equiv \frac{2|J^{++}| \cos(\arg J^{++}) \sin \kappa}{\mathcal{E}_{\kappa}}. \end{aligned} \quad (62)$$

The essential simplifications in expressions for contractions (59)-(62) take place in the case of model (30), that is when $J^{xy} = -J^{yx} = D$. Then

$$\begin{aligned}
\langle \varphi_j^+ \varphi_{j+n}^+ \rangle &= \delta_{n,0} + \frac{i}{N} \sum_{\kappa} \sin(\kappa n) \tanh \frac{\beta E_{\kappa}}{2} \equiv E(n), \\
\langle \varphi_j^- \varphi_{j+n}^- \rangle &= -E(n), \\
\langle \varphi_j^+ \varphi_{j+n}^- \rangle &= \frac{1}{N} \sum_{\kappa} \cos(\kappa n + \psi_{\kappa}) \tanh \frac{\beta E_{\kappa}}{2} \equiv G(n), \\
\langle \varphi_j^- \varphi_{j+n}^+ \rangle &= -G(-n),
\end{aligned} \tag{63}$$

where $\cos \psi_{\kappa} \equiv \epsilon_{\kappa}^{(+)} / \mathcal{E}_{\kappa}$, $\sin \psi_{\kappa} \equiv 2J^{++} \sin \kappa / \mathcal{E}_{\kappa}$.

Let's return to the evaluation of equal-time spin correlation functions and consider, for instance, $\langle s_j^x s_{j+n}^x \rangle$. For this correlation function with the utilization of (53)-(55) one derives

$$\langle s_j^x s_{j+n}^x \rangle = \frac{1}{4} \langle \varphi_j^- \varphi_{j+1}^+ \varphi_{j+1}^- \varphi_{j+2}^+ \cdots \varphi_{j+n-1}^+ \varphi_{j+n-1}^- \varphi_{j+n}^+ \rangle, \tag{64}$$

and after exploiting Wick-Bloch-de Dominicis theorem in r.h.s. of (64) for its square one gets the following expression

$$\begin{aligned}
& \left[4 \langle s_j^x s_{j+n}^x \rangle \right]^2 = \\
& \begin{vmatrix}
0 & -E(1) & \dots & -E(n-1) & -G(-1) & -G(-2) & \dots & -G(-n) \\
E(1) & 0 & \dots & -E(n-2) & -G(0) & -G(-1) & \dots & -G(-n+1) \\
E(2) & E(1) & \dots & -E(n-3) & -G(1) & -G(0) & \dots & -G(-n+2) \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
E(n-1) & E(n-2) & \dots & 0 & -G(n-2) & -G(n-3) & \dots & -G(-1) \\
G(-1) & G(0) & \dots & G(n-2) & 0 & E(1) & \dots & E(n-1) \\
G(-2) & G(-1) & \dots & G(n-3) & -E(1) & 0 & \dots & E(n-2) \\
G(-3) & G(-2) & \dots & G(n-4) & -E(2) & -E(1) & \dots & E(n-3) \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
G(-n) & G(-n+1) & \dots & G(-1) & -E(n-1) & -E(n-2) & \dots & 0
\end{vmatrix}.
\end{aligned} \tag{65}$$

In a similar way for other pair spin correlators one obtains

$$\left[4i \langle s_j^x s_{j+n}^y \rangle \right]^2 =$$

$$\begin{vmatrix}
0 & -E(1) & \dots -E(n-1) & -E(-n) & -G(-1) & \dots -G(-n+1) \\
E(1) & 0 & \dots -E(n-2) & -E(n-1) & -G(0) & \dots -G(-n+2) \\
E(2) & E(1) & \dots -E(n-3) & -E(n-2) & -G(1) & \dots -G(-n+3) \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
E(n-1) & E(n-2) & \dots & 0 & -E(1) & -G(n-2) \dots -G(0) \\
E(n) & E(n-1) & \dots & E(1) & 0 & -G(n-1) \dots -G(1) \\
G(-1) & G(0) & \dots & G(n-2) & G(n-1) & 0 \dots E(n-2) \\
G(-2) & G(-1) & \dots & G(n-3) & G(n-2) & -E(1) \dots E(n-3) \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
G(-n+1) & G(-n+2) \dots & G(0) & G(1) & -E(n-2) \dots & 0
\end{vmatrix}, \tag{66}$$

$$\langle s_j^x s_{j+n}^z \rangle = 0; \tag{67}$$

$$\left[4\ell \langle s_j^y s_{j+n}^x \rangle \right]^2 =
\begin{vmatrix}
0 & -E(1) & \dots -G(1) & -G(0) & -G(-1) & \dots -G(-n+1) \\
E(1) & 0 & \dots -G(2) & -G(1) & -G(0) & \dots -G(-n+2) \\
E(2) & E(1) & \dots -G(3) & -G(2) & -G(1) & \dots -G(-n+3) \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
G(1) & G(2) & \dots & 0 & E(1) & E(2) \dots E(n) \\
G(0) & G(1) & \dots -E(1) & 0 & E(1) & \dots E(n-1) \\
G(-1) & G(0) & \dots -E(2) & -E(1) & 0 & \dots E(n-2) \\
G(-2) & G(-1) & \dots -E(3) & -E(2) & -E(1) & \dots E(n-3) \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
G(-n+1) & G(-n+2) \dots -E(n) & -E(n-1) & -E(n-2) \dots & 0
\end{vmatrix}, \tag{68}$$

$$\left[4 \langle s_j^y s_{j+n}^y \rangle \right]^2 =$$

$$\begin{vmatrix}
0 & -E(1) & \dots & -E(n-1) & -G(1) & -G(0) & \dots & -G(-n+2) \\
E(1) & 0 & \dots & -E(n-2) & -G(2) & -G(1) & \dots & -G(-n+3) \\
E(2) & E(1) & \dots & -E(n-3) & -G(3) & -G(2) & \dots & -G(-n+4) \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
E(n-1) & E(n-2) & \dots & 0 & -G(n) & -G(n-1) & \dots & -G(1) \\
G(1) & G(2) & \dots & G(n) & 0 & E(1) & \dots & E(n-1) \\
G(0) & G(1) & \dots & G(n-1) & -E(1) & 0 & \dots & E(n-2) \\
G(-1) & G(0) & \dots & G(n-2) & -E(2) & -E(1) & \dots & E(n-3) \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
G(-n+2) & G(-n+3) & \dots & G(1) & -E(n-1) & -E(n-2) & \dots & 0
\end{vmatrix}, \quad (69)$$

$$\langle s_j^y s_{j+n}^z \rangle = 0; \quad (70)$$

$$\langle s_j^z s_{j+n}^x \rangle = 0, \quad (71)$$

$$\langle s_j^z s_{j+n}^y \rangle = 0, \quad (72)$$

$$\left[4 \langle s_j^z s_{j+n}^z \rangle \right]^2 = \begin{vmatrix}
0 & E(n) & G(0) & G(n) \\
-E(n) & 0 & G(-n) & G(0) \\
-G(0) & -G(-n) & 0 & -E(n) \\
-G(n) & -G(0) & E(n) & 0
\end{vmatrix}. \quad (73)$$

In figs. 9-12 the temperature dependences (1: $D = 0$, $\Omega = 0$, $1' : D = J^{xx}$, $\Omega = 0$; 2: $D = 0$, $\Omega = J^{xx}$, $2' : D = J^{xx}$, $\Omega = J^{xx}$) and in figs. 13-16 the dependences on transverse field (1: $D = 0$, $\beta = 1/J^{xx}$, $1' : D = J^{xx}$, $\beta = 1/J^{xx}$; 2: $D = 0$, $\beta = 10/J^{xx}$, $2' : D = J^{xx}$, $\beta = 10/J^{xx}$) for some pair spin static correlation functions are depicted. It is necessary to underline the peculiarities caused by the presence of Dzyaloshinskii-Moriya interaction. First, only $\langle s_j^x s_{j+n}^z \rangle$, $\langle s_j^y s_{j+n}^z \rangle$, $\langle s_j^z s_{j+n}^x \rangle$, $\langle s_j^z s_{j+n}^y \rangle$ are equal to zero, but not $\langle s_j^x s_{j+n}^y \rangle$ and $\langle s_j^y s_{j+n}^x \rangle$. The last two correlators tend to zero when $J^{xy} = J^{yx} = 0$. In this case $E(n) = 0$ for $n \neq 0$ and hence (66) and (68) may be rewritten as determinants of matrices with only non-zero rectangle (but not square) submatrices on their diagonals; such determinants are equal to zero. Second, the dependence of pair static correlation functions on n is nonmonotonic (in accordance with ref.[11] this fact indicates the appearance in the system of the incommensurate spiral spin structure).

5 Dynamics of transverse spin correlations and dynamical transverse susceptibility

Let's consider the dynamics of transverse spin correlations calculating for this purpose the transverse time-dependent (dynamical) pair spin correlation function $\langle s_j^z(t) s_{j+n}^z \rangle$. Due to the possibility of exploiting for its calculation c -cyclic Hamiltonian H^- (43) the evaluation of this correlation function in accordance with (53) and (52) reduces to estimation of dynamical correlation functions density-density for the system of non-interacting fermions

$$\begin{aligned}
4 \langle s_j^z(t) s_{j+n}^z \rangle &= \\
&= \sum_{\kappa_1, \kappa_2, \kappa_3, \kappa_4} \langle [\lambda_{j\kappa_1}^+ \beta_{\kappa_1}^+(t) + \mu_{j\kappa_1}^+ \beta_{-\kappa_1}(t)] [\lambda_{j\kappa_2}^- \beta_{\kappa_2}^+(t) - \mu_{j\kappa_2}^- \beta_{-\kappa_2}(t)] \times \\
&\quad \times [\lambda_{j+n, \kappa_3}^+ \beta_{\kappa_3}^+ + \mu_{j+n, \kappa_3}^+ \beta_{-\kappa_3}] [\lambda_{j+n, \kappa_4}^- \beta_{\kappa_4}^+ - \mu_{j+n, \kappa_4}^- \beta_{-\kappa_4}] \rangle = \\
&= \sum_{\kappa_1, \kappa_2, \kappa_3, \kappa_4} \left[-\lambda_{j\kappa_1}^+ \lambda_{j\kappa_2}^- \mu_{j+n, \kappa_3}^+ \mu_{j+n, \kappa_4}^- \langle \beta_{\kappa_1}^+ \beta_{\kappa_2}^+ \beta_{-\kappa_3} \beta_{-\kappa_4} \rangle e^{\imath(E_{\kappa_1} + E_{\kappa_2})t} + \right. \\
&\quad + \lambda_{j\kappa_1}^+ \mu_{j\kappa_2}^- \lambda_{j+n, \kappa_3}^+ \mu_{j+n, \kappa_4}^- \langle \beta_{\kappa_1}^+ \beta_{-\kappa_2} \beta_{\kappa_3}^+ \beta_{-\kappa_4} \rangle e^{\imath(E_{\kappa_1} - E_{-\kappa_2})t} - \\
&\quad - \lambda_{j\kappa_1}^+ \mu_{j\kappa_2}^- \mu_{j+n, \kappa_3}^+ \lambda_{j+n, \kappa_4}^- \langle \beta_{\kappa_1}^+ \beta_{-\kappa_2} \beta_{-\kappa_3} \beta_{\kappa_4}^+ \rangle e^{\imath(E_{\kappa_1} - E_{-\kappa_2})t} - \\
&\quad - \mu_{j\kappa_1}^+ \lambda_{j\kappa_2}^- \lambda_{j+n, \kappa_3}^+ \mu_{j+n, \kappa_4}^- \langle \beta_{-\kappa_1} \beta_{\kappa_2}^+ \beta_{\kappa_3}^+ \beta_{-\kappa_4} \rangle e^{\imath(E_{\kappa_2} - E_{-\kappa_1})t} + \\
&\quad + \mu_{j\kappa_1}^+ \lambda_{j\kappa_2}^- \mu_{j+n, \kappa_3}^+ \lambda_{j+n, \kappa_4}^- \langle \beta_{-\kappa_1} \beta_{\kappa_2}^+ \beta_{-\kappa_3} \beta_{\kappa_4}^+ \rangle e^{\imath(E_{\kappa_2} - E_{-\kappa_1})t} - \\
&\quad \left. - \mu_{j\kappa_1}^+ \mu_{j\kappa_2}^- \lambda_{j+n, \kappa_3}^+ \lambda_{j+n, \kappa_4}^- \langle \beta_{-\kappa_1} \beta_{-\kappa_2} \beta_{\kappa_3}^+ \beta_{\kappa_4}^+ \rangle e^{-\imath(E_{-\kappa_1} + E_{-\kappa_2})t} \right]. \quad (74)
\end{aligned}$$

In r.h.s. of (74) only non-zero averages of β -operators are written down and the following relations

$$\beta_{\kappa}^+(t) = \beta_{\kappa}^+ \exp(\imath E_{\kappa} t), \quad \beta_{\kappa}(t) = \beta_{\kappa} \exp(-\imath E_{\kappa} t) \quad (75)$$

were used. The averages of β -operators can be calculated using Wick-Bloch-de Dominicis theorem, e.g.

$$\begin{aligned}
\langle \beta_{\kappa_1}^+ \beta_{\kappa_2}^+ \beta_{-\kappa_3} \beta_{-\kappa_4} \rangle &= \\
&= -\frac{\delta_{\kappa_1, -\kappa_3}}{1 + e^{\beta E_{\kappa_1}}} \frac{\delta_{\kappa_2, -\kappa_4}}{1 + e^{\beta E_{\kappa_2}}} + \frac{\delta_{\kappa_1, -\kappa_4}}{1 + e^{\beta E_{\kappa_1}}} \frac{\delta_{\kappa_2, -\kappa_3}}{1 + e^{\beta E_{\kappa_2}}} = \\
&= -f_{\kappa_1} f_{\kappa_2} \delta_{\kappa_1, -\kappa_3} \delta_{\kappa_2, -\kappa_4} + f_{\kappa_1} f_{\kappa_2} \delta_{\kappa_1, -\kappa_4} \delta_{\kappa_2, -\kappa_3} \quad (76)
\end{aligned}$$

etc. After computation of these averages one finds that the coefficients near the averages contain the following products $\lambda_{j\kappa}^+ \mu_{j+n, -\kappa}^+$, $\lambda_{j+n, \kappa}^+ \mu_{j, -\kappa}^+$,

$\lambda_{j\kappa}^- \mu_{j+n,-\kappa}^-$, $\lambda_{j+n,\kappa}^- \mu_{j,-\kappa}^-$, $\lambda_{j\kappa}^+ \mu_{j+n,-\kappa}^-$, $\lambda_{j+n,\kappa}^+ \mu_{j,-\kappa}^-$, $\lambda_{j\kappa}^- \mu_{j+n,-\kappa}^+$, $\lambda_{j+n,\kappa}^- \mu_{j,-\kappa}^+$. They can be found with the help of (52). For simplicity in what follows their values will be used in the case when $J^{xy} = -J^{yx} = D$. Then

$$\begin{aligned}\lambda_{j\kappa}^+ \mu_{j+n,-\kappa}^+ &= \frac{1}{N} e^{-i\kappa n} = \lambda_{j\kappa}^- \mu_{j+n,-\kappa}^-, \\ \lambda_{j+n,\kappa}^+ \mu_{j,-\kappa}^+ &= \frac{1}{N} e^{i\kappa n} = \lambda_{j+n,\kappa}^- \mu_{j,-\kappa}^-, \\ \lambda_{j\kappa}^+ \mu_{j+n,-\kappa}^- &= \frac{1}{N} e^{-i(\kappa n + \psi_\kappa)}, \quad \lambda_{j+n,\kappa}^+ \mu_{j,-\kappa}^- = \frac{1}{N} e^{i(\kappa n - \psi_\kappa)}, \\ \lambda_{j\kappa}^- \mu_{j+n,-\kappa}^+ &= \frac{1}{N} e^{-i(\kappa n - \psi_\kappa)}, \quad \lambda_{j+n,\kappa}^- \mu_{j,-\kappa}^+ = \frac{1}{N} e^{i(\kappa n + \psi_\kappa)}.\end{aligned}\quad (77)$$

Gathering (74)-(77) together one derives the desired expression for transverse time-dependent correlation function for the model (30)

$$\begin{aligned}4 < s_j^z(t) s_{j+n}^z > = \\ &= \left[\frac{1}{N} \sum_{\kappa} \frac{\cosh \left(-iE_{\kappa} t + i\kappa n + \frac{\beta E_{\kappa}}{2} \right)}{\cosh \left(\frac{\beta E_{\kappa}}{2} \right)} \right]^2 + \left[\frac{1}{N} \sum_{\kappa} \frac{\sinh \left(i\psi_{\kappa} + \frac{\beta E_{\kappa}}{2} \right)}{\cosh \left(\frac{\beta E_{\kappa}}{2} \right)} \right]^2 - \\ &- \left[\frac{1}{N} \sum_{\kappa} \frac{\sinh \left(-iE_{\kappa} t + i\kappa n + i\psi_{\kappa} + \frac{\beta E_{\kappa}}{2} \right)}{\cosh \left(\frac{\beta E_{\kappa}}{2} \right)} \right] \times \\ &\times \left[\frac{1}{N} \sum_{\kappa} \frac{\sinh \left(-iE_{\kappa} t + i\kappa n - i\psi_{\kappa} + \frac{\beta E_{\kappa}}{2} \right)}{\cosh \left(\frac{\beta E_{\kappa}}{2} \right)} \right].\end{aligned}\quad (78)$$

Although E_{κ} and $\cos \psi_{\kappa}$, $\sin \psi_{\kappa}$ in (78) are determined by formulae (43), (63) for the case $J^{xy} = -J^{yx} = D$ the obtained result covers the case (26) as well. Remembering formulae (27) and (31) one should simply use $J^{xx} \cos^2 \alpha + \frac{J^{xy} + J^{yx}}{2} \sin 2\alpha + J^{yy} \sin^2 \alpha$ instead of J^x , $J^{xx} \sin^2 \alpha - \frac{J^{xy} + J^{yx}}{2} \sin 2\alpha + J^{yy} \cos^2 \alpha$ instead of J^y , and $\frac{J^{xy} - J^{yx}}{2}$ instead of D with $\tan 2\alpha = (J^{xy} + J^{yx}) / (J^{xx} - J^{yy})$. If one puts $D = 0$ in (78) it transforms into the well-known result obtained by Th.Niemeijer [21]. The depicted in figs. 17-20 dependence of the transverse dynamical autocorrelation function (78) on time ($\beta = 10/J^{xx}$; 1 : $D = 0$, $\Omega = 0$, 1' : $D = J^{xx}$, $\Omega = 0$; 2 : $D = 0$, $\Omega = J^{xx}$, 2' : $D = J^{xx}$, $\Omega = J^{xx}$) shows substantial changes caused by Dzyaloshinskii-Moriya interaction.

The dynamical susceptibility

$$\chi_{\alpha\beta}(\kappa, \omega) \equiv \sum_{n=1}^N e^{i\kappa n} \int_0^{\infty} dt e^{i(\omega + i\varepsilon)t} \frac{1}{i} < [s_j^{\alpha}(t), s_{j+n}^{\beta}] > \quad (79)$$

is of great interest from the point of view of observable properties of the system. The obtained result (78) permits one to calculate the transverse dynamical susceptibility. Really, taking into account the translation invariance one gets

$$\begin{aligned}
& \langle [s_j^z(t), s_{j+n}^z] \rangle = \langle s_j^z(t) s_{j+n}^z \rangle - \langle s_j^z(-t) s_{j-n}^z \rangle = \\
& = i \left[\frac{1}{N} \sum_{\kappa} \cos(\kappa n - E_{\kappa} t) \right] \left[\frac{1}{N} \sum_{\kappa} \sin(\kappa n - E_{\kappa} t) \tanh \frac{\beta E_{\kappa}}{2} \right] - \\
& - \frac{i}{2} \left\{ \left[\frac{1}{N} \sum_{\kappa} \cos(\kappa n - E_{\kappa} t - \psi_{\kappa}) \tanh \frac{\beta E_{\kappa}}{2} \right] \left[\frac{1}{N} \sum_{\kappa} \sin(\kappa n - E_{\kappa} t + \psi_{\kappa}) \right] + \right. \\
& \left. + \left[\frac{1}{N} \sum_{\kappa} \cos(\kappa n - E_{\kappa} t + \psi_{\kappa}) \tanh \frac{\beta E_{\kappa}}{2} \right] \left[\frac{1}{N} \sum_{\kappa} \sin(\kappa n - E_{\kappa} t - \psi_{\kappa}) \right] \right\}. \quad (80)
\end{aligned}$$

Using for summation over sites in (79) the lattice sum $\frac{1}{N} \sum_{n=1}^N e^{i\kappa n} = \delta_{\kappa,0}$, evaluating the integrals over t of the form

$$\int dt e^{i(\omega + F_{\kappa} + i\varepsilon)t} = \frac{i}{\omega + F_{\kappa} + i\varepsilon}, \quad (81)$$

bearing in mind the definition of functions $\cos \psi_{\kappa}$, $\sin \psi_{\kappa}$, and performing thermodynamical limit one ends up with

$$\begin{aligned}
\chi_{zz}(\kappa, \omega) = & \frac{1}{8\pi} \int_{-\pi}^{\pi} d\rho \left[\frac{1 + \cos(\psi_{\rho} + \psi_{\rho-\kappa})}{E_{\rho-\kappa} - E_{\rho} - \omega - i\varepsilon} + \frac{1 - \cos(\psi_{\rho} + \psi_{\rho-\kappa})}{-E_{\kappa-\rho} - E_{\rho} - \omega - i\varepsilon} - \right. \\
& \left. - \frac{1 + \cos(\psi_{\rho} + \psi_{\rho+\kappa})}{-E_{\rho+\kappa} + E_{\rho} - \omega - i\varepsilon} - \frac{1 - \cos(\psi_{\rho} + \psi_{\rho+\kappa})}{E_{-\rho-\kappa} + E_{\rho} - \omega - i\varepsilon} \right] \tanh \frac{\beta E_{\rho}}{2}. \quad (82)
\end{aligned}$$

Using the relation

$$\frac{1}{F_{\rho} - \omega - i\varepsilon} = \mathcal{P} \frac{1}{F_{\rho} - \omega} + i\pi \delta(F_{\rho} - \omega), \quad (83)$$

for real and imaginary parts of transverse susceptibility one gets final expressions

$$\begin{aligned}
Re \chi_{zz}(\kappa, \omega) = & \frac{1}{8\pi} \mathcal{P} \int_{-\pi}^{\pi} d\rho \left[\frac{1 + \cos(\psi_{\rho} + \psi_{\rho-\kappa})}{E_{\rho-\kappa} - E_{\rho} - \omega} + \frac{1 - \cos(\psi_{\rho} + \psi_{\rho-\kappa})}{-E_{\kappa-\rho} - E_{\rho} - \omega} - \right. \\
& \left. - \frac{1 + \cos(\psi_{\rho} + \psi_{\rho+\kappa})}{-E_{\rho+\kappa} + E_{\rho} - \omega} - \frac{1 - \cos(\psi_{\rho} + \psi_{\rho+\kappa})}{E_{-\rho-\kappa} + E_{\rho} - \omega} \right] \tanh \frac{\beta E_{\rho}}{2}, \quad (84)
\end{aligned}$$

$$\begin{aligned}
Im\chi_{zz}(\kappa, \omega) = & \frac{1}{8} \int_{-\pi}^{\pi} d\rho \{ [1 + \cos(\psi_{\rho} + \psi_{\rho-\kappa})] \delta(E_{\rho-\kappa} - E_{\rho} - \omega) + \\
& + [1 - \cos(\psi_{\rho} + \psi_{\rho-\kappa})] \delta(-E_{\kappa-\rho} - E_{\rho} - \omega) - \\
& - [1 + \cos(\psi_{\rho} + \psi_{\rho+\kappa})] \delta(-E_{\rho+\kappa} + E_{\rho} - \omega) - \\
& - [1 - \cos(\psi_{\rho} + \psi_{\rho+\kappa})] \delta(E_{-\rho-\kappa} + E_{\rho} - \omega) \} \tanh \frac{\beta E_{\rho}}{2}. \quad (85)
\end{aligned}$$

These are the main results of the present paper.

It is useful to look at the particular case $\kappa = 0$. In this case one has

$$\begin{aligned}
\chi_{zz}(0, \omega) = & \frac{1}{4\pi} \int_{-\pi}^{\pi} d\rho \sin^2 \psi_{\rho} \left[\frac{1}{-E_{\rho} - E_{-\rho} - \omega - i\varepsilon} - \right. \\
& \left. - \frac{1}{E_{\rho} + E_{-\rho} - \omega - i\varepsilon} \right] \tanh \frac{\beta E_{\rho}}{2} \quad (86)
\end{aligned}$$

so that, for instance,

$$Im\chi_{zz}(0, \omega) = -\frac{1}{4} \int_{-\pi}^{\pi} d\rho \sin^2 \psi_{\rho} \delta(2\mathcal{E}_{\rho} - \omega) \tanh \frac{\beta E_{\rho}}{2}. \quad (87)$$

In the case of isotropic XY model with Dzyaloshinskii-Moriya interaction $\sin \psi_{\rho} = 0$ and $Im\chi_{zz}(0, \omega) = 0$ as one should expect because in this case $[\sum_{j=1}^N s_j^z, H] = 0$. In the case of de Gennes model with Dzyaloshinskii-Moriya interaction when $\mathcal{E}_{\rho} = \sqrt{\Omega^2 + \Omega J \cos \rho + J^2/4}$ one can integrate in (87) over ρ using the relation

$$\delta(2\mathcal{E}_{\rho} - \omega) = \sum_{\rho_0} \frac{\delta(\rho - \rho_0)}{2 \left| \frac{\partial \mathcal{E}_{\rho}}{\partial \rho} \right|}, \quad (88)$$

where by $\rho_0 = \rho_0(\omega)$ the solutions of the equation $2\mathcal{E}_{\rho_0} - \omega = 0$ are denoted. This equation can be written in the form

$$\cos \rho_0 = \frac{\omega^2 - J^2 - 4\Omega^2}{4\Omega J}, \quad (89)$$

and when ω satisfies inequalities

$$-1 \leq \frac{\omega^2 - J^2 - 4\Omega^2}{4\Omega J} \leq 1 \quad (90)$$

or for $\Omega, J > 0$

$$|J - 2\Omega| \leq \omega \leq J + 2\Omega, \quad (91)$$

equation (89) has two solutions in the interval of integration $\rho_0 \geq 0$ and $-\rho_0$. Besides $\partial \mathcal{E}_\rho / \partial \rho = -\Omega J \sin \rho / 2\mathcal{E}_\rho$, $\sin \psi_\rho = J \sin \rho / 2\mathcal{E}_\rho$, so that in the case of de Gennes model with Dzyaloshinskii-Moriya interaction one gets the following final result

$$\begin{aligned} \text{Im} \chi_{zz}(0, \omega) = \\ = \begin{cases} \frac{-J |\sin \rho_0|}{16 \mathcal{E}_{\rho_0} \Omega} \left(\tanh \frac{\beta E_{\rho_0}}{2} + \tanh \frac{\beta E_{-\rho_0}}{2} \right), & \text{if } |J - 2\Omega| \leq \omega \leq J + 2\Omega, \\ 0, & \text{otherwise.} \end{cases} \end{aligned} \quad (92)$$

The presented in figs. 21,22 results of the numerical calculations of frequency dependence of $\text{Im} \chi_{zz}(0, \omega)$ (86) for de Gennes model with Dzyaloshinskii-Moriya interaction ($\beta = 10/J^{xx}$; 1 : $D = 0$, 2 : $D = 0.5J^{xx}$, 3: $D = J^{xx}$) show that the presence of this interaction dramatically changes the frequency dependence. This fact seems to be of great importance in connection with the possible experimental prove of the presence in the system of Dzyaloshinskii-Moriya interaction on the base of experimental measurements of $\text{Im} \chi_{zz}(0, \omega)$.

6 Conclusions

Let's sum up the results of present study of statistical mechanics of 1D $s = \frac{1}{2}$ XY anisotropic ring in transverse field with Dzyaloshinskii-Moriya interaction. This interaction keeps the model in the class of 1D $s = \frac{1}{2}$ XY models because after fermionization of the Hamiltonian one is faced with the quadratic in Fermi operators forms. However, after their diagonalization one finds that the spectrum E_κ no longer is even function of κ . This leads only to some technical complications in computations. The obtained thermodynamical functions and static spin correlation functions essentially depend on the value of Dzyaloshinskii-Moriya interaction. For instance, these interaction decreases the transverse magnetization at certain transverse field in de Gennes model and in isotropic XY model (figs. 5,6). They lead to appearance of non-zero spin correlators $\langle s_j^x s_{j+n}^y \rangle$ and $\langle s_j^y s_{j+n}^x \rangle$ and to nonmonotonic dependence of pair spin correlation functions on n . The evaluation of transverse dynamical correlation function and the corresponding susceptibility shows that Dzyaloshinskii-Moriya interaction essentially influences on the dynamics of transverse spin correlations (figs. 17-20) and drastically changes the dynamical susceptibility (figs. 21,22).

In addition it is necessary to note that if $J^{xy} = J^{yx} = 0$ all obtained results transform into the corresponding results for anisotropic XY model.

Really, in this limit $E_\kappa \rightarrow \mathcal{E}_\kappa \rightarrow \sqrt{\epsilon_\kappa^{(+)^2} + 4(J^{++})^2 \sin^2 \kappa}$ and $E_\kappa = E_{-\kappa}$. Due to this simplification the thermodynamical functions because of parity of integrands contain $2 \int_0^\pi d\kappa(\dots)$ instead of $\int_{-\pi}^\pi d\kappa(\dots)$. In contractions (59)-(63) $\langle \varphi_j^+ \varphi_{j+n}^+ \rangle \rightarrow \delta_{n,0}$, $\langle \varphi_j^- \varphi_{j+n}^- \rangle \rightarrow -\delta_{n,0}$, $\langle \varphi_j^+ \varphi_{j+n}^- \rangle \rightarrow \frac{1}{\pi} \int_0^\pi d\kappa \cos(\kappa n + \psi_\kappa) \tanh \frac{\beta E_\kappa}{2} \equiv G(n)$, $\langle \varphi_j^- \varphi_{j+n}^+ \rangle \rightarrow -G(-n)$ so that $4 \langle s_j^x s_{j+n}^x \rangle$, $4 \langle s_j^y s_{j+n}^y \rangle$, $4 \langle s_j^z s_{j+n}^z \rangle$ (but not their squares) can be rewritten as $N \times N$ determinants and $\langle s_j^x s_{j+n}^y \rangle = \langle s_j^y s_{j+n}^x \rangle = 0$. The transverse dynamical correlation function transforms into the obtained in ref.[21] expression.

The performed investigations follow earlier works [18,22-29] considering the derivation of exact results in statistical mechanics of 1D $s = \frac{1}{2}$ systems with Dzyaloshinskii-Moriya interaction.

At last it should be mentioned that for a quite a lot of magnetic and ferroelectric materials, showing nearly 1D behavior above their ordering temperatures, a variety of experimental data are now available [30-40] and thus theoretical investigations of statistical mechanics of 1D spin models may be of great interest for clarifying whether the properties of such simple spin models are capable to caricature the measurements.

The authors would like to thank J.Rossat-Mignod, F.P.Onufrieva and other participants of the Ukrainian-French Symposium "Condensed Matter: Science & Industry" (Lviv, 20-27 February 1993) for stimulating discussions. They would like to express the gratitude to the participants of the seminar of Quantum Statistics Department of ICMP (11.05.1993) and to the participants of seminars of Laboratory for the Theory of Model Spin Systems of this department for many helpful discussions.

References

- [1] Lieb E., Schultz T., Mattis D. Two soluble models of an antiferromagnetic chain//Annals of Physics.-1961. -**16**,3.-P.407-466.
- [2] Katsura S. Statistical mechanics of the anisotropic linear Heisenberg model//Phys.Rev.-1962.-**127**,5.-P.1508-1518.
- [3] Chesnut D.B., Suna A. Fermion behavior of one-dimensional excitons//J.Chem.Phys.-1963.-**39**,1.-P.146-149.

- [4] Pikin S.A., Tsukernik V.M. Thermodynamics of linear spin chains in a transverse magnetic field//Zh.Eksp.Teor.Fiz.-1966.- **50**,5.-P.1377-1380.(in Russian)
- [5] Mazur P., Siskens Th.J. Time correlation functions in the a -cyclic XY model. I//Physica.-1973.-**69**.-P.259-272.
- [6] Siskens Th.J., Mazur P. Time-correlation functions in the a -cyclic XY model. II//Physica.-1974.-**71**.-P.560-578.
- [7] Goncalves L.L. Dynamics of the one-dimensional transverse Ising model//J.Phys.A.-1980.-**13**.-P.223-236.
- [8] McCoy B.M., Barouch E., Abraham D.B. Statistical mechanics of the XY model. IV. Time-dependent spin-correlation functions // Phys.Rev.A.-1971.-**4**,6.-P.2331-2342.
- [9] Barouch E., Dresden M. Exact time-dependent analysis for the one-dimensional XY model // Phys.Rev.Lett.-1969.-**23**,3.-P.114-117.
- [10] Nishimori H. One-dimensional XY model in Lorentzian random field // Phys.Lett.A.-1984.-**100**,5.-P.239-243.
- [11] Kontorovich V.M., Tsukernik V.M. Spiral structure in the one-dimensional spin chain // Zh.Eksp.Teor.Fiz.-1967.-**52**,5.-P.1446-1453. (in Russian)
- [12] Dzyaloshinsky I.E. A thermodynamical theory of "weak" ferromagnetism in antiferromagnetic substances//Zh.Eksp.Teor.Fiz.-1957.-**32**,6.-P.1547-1562. (in Russian)
- [13] Moriya T. Anisotropic superexchange interaction and weak ferromagnetism//Phys.Rev.-1960.-**120**,1.-P.91-98.
- [14] Anderson P.W. New approach to the theory of superexchange interactions//Phys.Rev.-1959.-**115**,1.-P.2-13.
- [15] Messia A. Kvantovaya mehanika, tom 2.-Moskva: Nauka, 1979.-584 p. (in Russian)
- [16] Izyumov Yu.A., Ozerov R.P. Magnitnaya neitronografiya.-Moskva: Nauka, 1966.-532 p. (in Russian)

- [17] Izyumov Yu.A. Difraktsiya neutronov na dlinnoperiodicheskikh strukturah.-Moskva: Energoatomizdat, 1987.-200 p. (in Russian)
- [18] Derzhko O.V., Levitskii R.R., Moina A.Ph. Approximation of Bose commutation rules and elementary excitation spectrum in the spin systems theory//Fiz.Kond.Sys.-1993.-1.-P.115-118. (in Ukrainian)
- [19] Isihara A. Statistical physics.- New York-London: Academic Press, 1971.
- [20] Baxter R.J. Exactly solved models in statistical mechanics.- London New York Paris San Diego San Francisco Sao Paulo Sydney Tokyo Toronto: Academic Press, 1982.
- [21] Niemeijer Th. Some exact calculations on a chain of spins $\frac{1}{2}$ //Physica.-1967.-**36**,3.-P.377-419.
- [22] Krivoruchko V.N. Magnon bound states in an anisotropic chain of spins with the Dzyaloshinskii interaction//Fiz.Niz.Temp.-1986.-**12**,8.- P.872-876. (in Russian)
- [23] Zvyagin A.A. The ground state structure of a spin chain with the Dzyaloshinsky type interaction//Fiz.Niz.Temp.-1989.-**15**,9.-P.977-979. (in Russian)
- [24] Zvyagin A.A. Characteristics of two-sublattice spin chain with coupling of Dzyaloshinsky type //Zh.Eksp.Teor.Fiz.-1990.-**98**,4.-P.1396-1401. (in Russian)
- [25] Zvyagin A.A. The effect of impurities on planar structure of spin chain with Dzyaloshinsky interaction//Fiz.Niz.Temp.-1991.-**17**,1.- P.125-127. (in Russian)
- [26] Zvyagin A.A. The transverse structure of a spin chain with Dzyaloshinskii-Moriya-type interaction //J.Phys.: Condens.Matter.-1991.-**3**.- P.3865-3867.
- [27] Derzhko O.V., Smereka I.T., Moina A.Ph. Thermodynamical and dynamical properties of XY-type 1D $s = \frac{1}{2}$ -spin models and some applications in condensed matter physics //Ukrainian-French Symposium "Condensed Matter: Science & Industry" (Lviv, February, 20-27, 1993). Abstracts, information & participants.-Lviv,1993.-P.198.

- [28] Derzhko O.V., Moina A.Ph. 1D $s = \frac{1}{2}$ anisotropic XY model in transverse field with Dzyaloshinskii-Moriya interaction //The Eighth International Meeting on Ferroelectricity, 8-13 August 1993. Program Summary and Abstract Book. Gaithersburg, Maryland, U.S.A.-P.80.
- [29] Derzhko O.V., Moina A.Ph., Levitskii R.R. Bose commutation rules approximation in spin systems theory//The Eighth International Meeting on Ferroelectricity, 8-13 August 1993. Program Summary and Abstract Book. Gaithersburg, Maryland, U.S.A.-P.80.
- [30] Steiner M., Villain J., Windsor C.G. Theoretical and experimental studies on one-dimensional magnetic systems//Advances in Physics.-1976.-**25**, 2.-P.87-209.
- [31] Mikeska H.-J., Steiner M. Solitary excitations in one-dimensional magnets //Advances in Physics.-1991.-**40**,3.-P.191-356.
- [32] Landee C.P., Willett R.D. Tetramethylammonium copper chloride and *tris* (trimethylammonium) copper chloride: $s = \frac{1}{2}$ Heisenberg one-dimensional ferromagnets//Phys.Rev.Lett.-1979.-**43**, 6.-P.463-466.
- [33] Willett R.D., Landee C.P., Gaura R.M., Swank D.D., Groenedijk H.A., van Duynveldt A.J. Magnetic properties of one-dimensional spin $\frac{1}{2}$ ferromagnets: metamagnetic behavior of $(C_6H_{11}NH_3)CuCl_3$ // Journal of Magnetism and Magnetic Materials.-1980.-**15-18**.-P.1055-1056.
- [34] Takahashi M., Turek Ph., Nakazawa Y., Tamura M., Nozawa K., Shiomori D., Ishikawa M., Kinoshita M. Discovery of a quasi-1D organic ferromagnet, *p*-NPNN//Phys.Rev.Lett.-1991.-**67**,6.-P.746-748.
- [35] Silverstein A.J., Soos Z.G. Comparison of three self-consistent ground states for the linear Heisenberg antiferromagnet//J.Chem.Phys.-1970.-**53**,1.-P.326-333.
- [36] De Carvalho A.V., Salinas S.R. Theory of phase transition in the quasi-one-dimensional hydrogen-bonded ferroelectric crystal $PbHPO_4$ // J.Phys.Soc.Japan.-1978.-**44**,1.-P.238-243.
- [37] Zinenko V.I. Theory of ferroelectrics of $PbHPO_4$ type //Fiz.Tv.Tela.-1979.-**21**,6.-P.1819-1825. (in Russian)

- [38] Levitsky R.R., Grigas J., Zachek I.R., Mits Ye.V., Paprotny W. Relaxational dynamics of quasi-one-dimensional CsD_2PO_4 -type ferroelectrics//Ferroelectrics.-1986.-**67**.-P.109-124.
- [39] D'Iorio M., Armstrong R.L., Taylor D.R. Longitudinal and transverse spin dynamics of a one-dimensional XY system studied by chlorine nuclear relaxation in $PrCl_3$ //Phys.Rev.B.-1983.-**27**,3.-P.1664-1673.
- [40] D'Iorio M., Glaus U., Stoll E. Transverse spin dynamics of a one-dimensional XY system: a fit to spin-spin relaxation data//Solid State Communications.-1983.-**47**,5.-P.313-315.

Figures caption

- Fig.1. $E_\kappa/J^{xx} = (\epsilon_\kappa^{(-)} + \mathcal{E}_\kappa)/J^{xx}$ vs. κ ; $J^{yy} = 0$.
- Fig.2. $E_\kappa/J^{xx} = (\epsilon_\kappa^{(-)} + \mathcal{E}_\kappa)/J^{xx}$ vs. κ ; $J^{yy} = J^{xx}$.
- Fig.3. c vs. $1/(\beta J^{xx})$; $J^{yy} = 0$.
- Fig.4. c vs. $1/(\beta J^{xx})$; $J^{yy} = J^{xx}$.
- Fig.5. $-\langle \frac{1}{N} \sum_{j=1}^N s_j^z \rangle$ vs. Ω/J^{xx} ; $J^{yy} = 0$.
- Fig.6. $-\langle \frac{1}{N} \sum_{j=1}^N s_j^z \rangle$ vs. Ω/J^{xx} ; $J^{yy} = J^{xx}$.
- Fig.7. $-\langle \frac{1}{N} \sum_{j=1}^N s_j^z \rangle$ vs. $1/(\beta J^{xx})$; $J^{yy} = 0$.
- Fig.8. $-\langle \frac{1}{N} \sum_{j=1}^N s_j^z \rangle$ vs. $1/(\beta J^{xx})$; $J^{yy} = J^{xx}$.
- Fig.9. $16 < s_j^x s_{j+1}^x >^2$ vs. $1/(\beta J^{xx})$; $J^{yy} = 0$.
- Fig.10. $16 < s_j^x s_{j+1}^x >^2$ vs. $1/(\beta J^{xx})$; $J^{yy} = J^{xx}$.
- Fig.11. $16 < s_j^x s_{j+1}^y >^2$ vs. $1/(\beta J^{xx})$; $J^{yy} = 0$.

Fig.12. $16 \langle s_j^x s_{j+1}^y \rangle^2$ vs. $1/(\beta J^{xx})$; $J^{yy} = J^{xx}$.

Fig.13. $16 \langle s_j^x s_{j+1}^x \rangle^2$ vs. Ω/J^{xx} ; $J^{yy} = 0$.

Fig.14. $16 \langle s_j^x s_{j+1}^x \rangle^2$ vs. Ω/J^{xx} ; $J^{yy} = J^{xx}$.

Fig.15. $16 \langle s_j^x s_{j+1}^y \rangle^2$ vs. Ω/J^{xx} ; $J^{yy} = 0$.

Fig.16. $16 \langle s_j^x s_{j+1}^y \rangle^2$ vs. Ω/J^{xx} ; $J^{yy} = J^{xx}$.

Fig.17. $Re \langle s_j^z(t) s_j^z \rangle$ vs. $J^{xx}t$; $J^{yy} = 0$.

Fig.18. $Im \langle s_j^z(t) s_j^z \rangle$ vs. $J^{xx}t$; $J^{yy} = 0$.

Fig.19. $Re \langle s_j^z(t) s_j^z \rangle$ vs. $J^{xx}t$; $J^{yy} = J^{xx}$.

Fig.20. $Im \langle s_j^z(t) s_j^z \rangle$ vs. $J^{xx}t$; $J^{yy} = J^{xx}$.

Fig.21. $-Im \chi_{zz}(0, \omega)$ vs. ω/J^{xx} ; $\Omega/J^{xx} = 0.25$.

Fig.22. $-Im \chi_{zz}(0, \omega)$ vs. ω/J^{xx} ; $\Omega/J^{xx} = 0.5$.

Institute for Condensed Matter Physics
1 Svientsitskii Str
Lviv 290011
Ukraine
e-mail: icmp@sigma.icmp.lviv.ua